

ORIGINAL RESEARCH

Open Access



# Synthesis, characterization and stress-testing of a robust quillaja saponin stabilized oil-in-water phytocannabinoid nanoemulsion

Abhinandan Banerjee<sup>1\*</sup>, Justin Binder<sup>2</sup>, Rayan Salama<sup>1</sup> and John F. Trant<sup>1\*</sup> 

## Abstract

**Background:** This study describes the design, optimization, and stress-testing of a novel phytocannabinoid nanoemulsion generated using high-pressure homogenization. QNaturale<sup>®</sup>, a plant-derived commercial emulsifier containing quillaja saponin, was used to stabilize the lipid phase droplets in water. Stress-testing was performed on this nanoemulsion in order to evaluate its chemical and colloidal stability under the influence of different environmental factors, encompassing both physical and chemical stressors.

**Methods:** Extensive optimization studies were conducted to arrive at an ideal nanoemulsion formulation. A coarse emulsion containing 16.6 wt% CBD-enriched cannabis distillate and 83.4 wt% carrier (soybean) oil dispersed in 10 wt% QNaturale<sup>®</sup> (1.5 wt% quillaja saponin) solution after 10 homogenization cycles at a pressure of 30,000 psi produced a stable nanoemulsion. This nanoemulsion was then subjected to the stress studies.

**Results:** The optimized nanoemulsion had an average droplet diameter of *ca.* 120 nm and average droplet surface  $\zeta$  potentials of *ca.* -30 mV. It was imaged and characterized by a variety of protocols. It proved to be stable to droplet agglomeration and phase separation upon storage under ambient conditions for 6 weeks, as well as under a variety of physical stressors such as heat, cold, dilution, and carbonation. pH values  $\leq 2$  and moderately high salt concentrations ( $> 100$  mM), however, destabilized the nanoemulsion, eventually leading to phase separation. Cannabis potency, determined by HPLC, was detrimentally affected by any changes in the nanoemulsion phase stability.

**Conclusions:** Quillaja saponin stabilized cannabidiol(CBD)-enriched nanoemulsions are stable, robust systems even at low emulsifier concentrations, and are therefore significant from both a scientific as well as a commercial perspective.

**Keywords:** Cannabis, Nanoemulsion, High pressure homogenization, Stress testing

## Background

As the therapeutic benefits of cannabis become scientifically established, and the use of nutraceutical and recreational cannabis products is decriminalized in countries such as Argentina, Belgium, Georgia, Uruguay, South

Africa, etc., cannabis research is expected to flourish in the new decade (McIver 2017). Canada, especially, is expected to play a pioneering role in cannabis research, given that it is the only G7 country to have completely legalized medical as well as recreational cannabis products, whilst simultaneously creating a robust regulatory system for the purposes of quality assurance and safe dissemination of cannabis-containing consumer products (Government of Canada 2018).

\*Correspondence: [arb@uwindsor.ca](mailto:arb@uwindsor.ca); [john.trant@uwindsor.ca](mailto:john.trant@uwindsor.ca)

<sup>1</sup>Department of Chemistry and Biochemistry, University of Windsor, 401 Sunset Ave., Windsor, ON, Canada

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Cannabis is a highly complex mixture, consisting of over 120 phytocannabinoids with related chemical structures, along with a variety of terpenes and flavonoids (Turner et al. 2017). As many of these natural products interact with the central nervous system (CNS) receptors (Zou and Kumar 2018), cannabis could prove to be an exceedingly promising source for pharmacophore discovery. Amongst the phytocannabinoids,  $\Delta^9$ -tetrahydrocannabinol (THC) and cannabidiol (CBD) are by far the most abundant, and consequently, the most thoroughly examined (Hanus et al. 2016; Reekie et al. 2017). THC and CBD have demonstrated neuroprotective, immunomodulatory, as well as anti-inflammatory effects (Cameron and Hemingway 2020), leading to their inclusion as adjunctive treatment for malignant brain tumors, Parkinson's disease, Alzheimer's disease, multiple sclerosis, neuropathic pain, and childhood seizure disorders (Maroon and Bost 2018). Experimental studies are being conducted in order to examine anecdotal and preliminary scientific evidence of their benefits in alleviating psychiatric and mood disorders, such as schizophrenia, anxiety, depression, addiction, and post-traumatic stress disorder (Shahbazi et al. 2020; Scherma et al. 2018). While the psychoactivity of THC can be less desirable in some clinical settings, the non-psychoactive CBD, with its anti-inflammatory, anti-convulsive, and anti-emetic effects, is a prime candidate for the development of functional cannabinoid-based nutraceuticals (Parker et al. 2002; Khan et al. 2020). Other promising biological effects of cannabinoids include anti-obesity and antidiabetic effects of tetrahydrocannabivarin (THCV) (Abioye et al. 2020) and in developing fluoride- and alcohol-free anti-bacterial mouthwash and oral care products (Vasudevan and Stahl 2020).

CBD is highly lipophilic and hydrophobic, thereby making oral delivery routes for CBD an essential but under-explored field of research (Grotenhermen 2003). CBD has low oral bioavailability (*ca.* 6–9%) owing to its low aqueous solubility and its susceptibility to ready clearance during first pass metabolism; therefore, it is up to the formulation science community to devise new routes for orally administering cannabinoids to both increase and ensure reproducibility of blood plasma concentrations (Elgart et al. 2012). Protocols such as self-emulsifying drug delivery systems have been applied in the past to increase solubility and reduce first pass metabolism of lipophilic bioactives such as cannabinoids in the context of pharmaceutical preparations (Cherniakov et al. 2017).

A potential approach used to augment the bioaccessibility of orally administered lipophilic nutraceuticals is their incorporation into nanoemulsions (Jaiswal et al. 2015; Solans et al. 2005). Due to reduced droplet sizes and lower emulsifier concentrations in these systems relative to microemulsions or liposomes, loading of phytocannabinoids in the lipid phase of nanoemulsions may improve

drug bioaccessibility (Acosta 2009), prevent oxidative degradation of the cannabinoids (Sun et al. 2015), and provide a more palatable and “label friendly” cannabinoid ingestible (Raikos and Ranawana 2017).

A stable nanoemulsion has a bulk phase (here, water); a diffused phase (here, a solution of cannabis extract in an edible oil); and an emulsifier. The emulsifier is of paramount importance: not only does it prevent the lipid phase droplets from agglomerating and undergoing phase-separation, but it also dictates the taste and ‘mouth-feel’ of the resultant nanoemulsion (Mouritsen and Styrbaek 2017; Panghal et al. 2019). Excess emulsifier may very well imbue an emulsion with a bitter ‘chemical’ or metallic taste, or an unpleasant texture; moreover, cost, safety and regulatory issues also come into play. Naturally occurring, ‘label-friendly’ emulsifiers are therefore preferred when devising such nanoformulations. QNaturale<sup>®</sup> is a commercially available emulsifier containing quillaja saponin obtained from the bark of the *quillaja saponaria* Molina tree. It has previously been found to be effective for the fabrication of nanoemulsions by high-energy methods such as microfluidization (Bai et al. 2016). Structurally, quillaja saponin might best be described as a mixture of triterpenoid saponins comprising a hydrophobic quillaic acid backbone glycosylated with hydrophilic sugar moieties (Fig. 1). The interfacially active saponins in quillaja extracts consist of two hydrophilic sugar chains and one hydrophobic aglycone, generating a three-unit structure with distinct hydrophilic–hydrophobic–hydrophilic properties (Reichert et al. 2019). The emulsion-stabilizing properties of QNaturale<sup>®</sup> are attributed to strong electrostatic repulsion in quillaja saponin coated lipid droplets with high negative surface  $\zeta$ -potentials between pH 3 and 9 (Kharat and McClements 2019; Zhu et al. 2019). Additionally, the fast adsorption kinetics of QNaturale<sup>®</sup> onto lipid droplet surfaces also promotes the formation of stable quillaja saponin oil-in-water nanoemulsions with sub- $\mu\text{m}$  droplet sizes (Bai et al. 2016; Zhang et al. 2016). Another important component is the carrier oil, which modulates emulsion stability and bioactive bioaccessibility. For the purposes of this study, soybean oil, a biodegradable, non-toxic, and sustainably sourced edible oil was selected owing to its excellent miscibility with cannabis extracts (Hsu and Nacu 2003). Soybean oil and soy protein isolates have been used in the past as carrier media for pharmaceuticals (Takino et al. 1994; Chung et al. 2001) and nutraceuticals (Teng et al. 2012; Chen and Subirade 2009).

This report adds to the few published accounts of cannabinoid emulsions as potential medicinal and/or nutraceutical products. Recently, Marangoni et al. reviewed THC encapsulation and stability in food matrices (Marangoni and Marangoni 2019); similarly, McClements has reviewed scientific developments in the



of the following cannabinoids was obtained from Cerilliant: cannabidiolic acid (CBD-A), tetrahydrocannabinolic acid (THC-A), cannabidiol (CBD), cannabigerolic acid (CBG-A), cannabigerol (CBG), cannabichromene (CBC), cannabinol (CBN), and delta-9 tetrahydrocannabinol (THC).

### Preparation of the nanoemulsions

A representative nanoemulsion was synthesized as follows: the CD<sub>CBD</sub> (69% CBD; see Table S1) was magnetically stirred and heated in a water bath at *ca.* 65°C, and to it was added the requisite weight of the soybean oil in order to create a homogeneous lipid phase. The stirring was continued for another 10–15 min. Meanwhile, QNaturale<sup>®</sup> was dissolved in water in order to generate the aqueous phase. Finally, the lipid phase was added to the water phase in the correct proportion (usually, 10 wt% lipid phase in water) while subjecting the mixture to three cycles of blending using a high shear mixer (IKA-T-1000) at ambient temperatures (60 s on, 60 s off at power setting = 3). This generated a milky coarse emulsion, to which was added 0.1 wt% potassium sorbate, and enough citric acid to lower the pH to ~3.4–3.8. The nanoemulsions were then formed by passing the coarse emulsions through a high pressure homogenizer (Nano DeBee, BEE International, USA; equipped with a chill loop) at a pressure of 30000 psi. The number of passes were optimized during the course of the study (*vide infra*). Finally, the nanoemulsions were poured in glass vials, capped, and stored either in a fridge at ~4°C in amber vials (for long term storage) or under ambient conditions in the dark (for short-term storage). The CBD concentration of the final emulsion was determined to be 10–11 mg.g<sup>-1</sup> (slight variations are attributed to individual lots of emulsions made) on the basis of measured weights, although quantitative analysis of the nanoemulsion revealed slight loss of CBD during the emulsion preparation (*vide infra*).

### Emulsion droplet size distributions

In order to determine lipid particle size distributions in the nanoemulsions, dynamic light scattering (DLS), also known as photon correlation spectroscopy, was applied. This technique generated Z-average diameters of the dispersed phase droplets ( $d_z$ ), as well as the polydispersity index (PDI). Diluted samples (50- to 100-fold dilutions) were used to avoid multiple scattering. The measurements were conducted with the Zetasizer (Nano ZS, Malvern Instruments Ltd., UK).  $d_z$  was calculated from the auto-correlation function of the intensity of light scattered from the particles. The software used was Zetasizer Software 7.13, supplied by the manufacturer (Malvern Instruments Ltd.) Disposable poly(styrene) cuvettes (ZEN0040) were used for sample measurements. Phase-separated

nanoemulsions were re-mixed by shaking prior to dilution for DLS measurements.

### Emulsion droplet zeta potential distributions

The rate of droplet movement under the influence of an external oscillating electrical field with a voltage of 150 V (electrophoretic mobility) was measured with the Zetasizer (Nano ZS, Malvern Instruments Ltd., UK) in folded capillary zeta-cells obtained from Malvern (DTS1070). The measured electrophoretic mobilities were converted to  $\zeta$ -potentials by the instrument software (Zetasizer Software 7.13, Malvern Instruments Ltd., UK) using Henry's equation:

$$U_e = \frac{2\varepsilon\zeta}{3\eta} f_{ak} \quad (1)$$

where  $U_e$  is the electrophoretic mobility,  $\varepsilon$  is the dielectric constant,  $\zeta$  is the zeta potential,  $\eta$  is the viscosity of the dispersant, and  $f_{ak}$  is the Henry function. The Smoluchowski approximation,  $f_{ak} = 1.5$ , was used for high ionic strength media, given that water was the bulk phase in all the measured emulsions (Sze et al. 2003).

### Optical microscopy

For optical microscopy, 10  $\mu$ L of the nanoemulsion was placed on a clean glass slide. A glass cover-slip was then positioned upon the droplet, and sealed in place with transparent nail-polish. Samples were examined and photographed immediately under an Olympus BX-51 microscope equipped with an Olympus U-CAMD3 camera under bright-field conditions at a magnification of 50x.

### Scanning electron microscopy (SEM)

SEM was performed on an FEI Nova NanoSEM430. Nanoemulsion droplets were dispersed on polished silicon (100) chips and dried, followed by sputter coating to 5–6 nm with iridium using a Lecai EM ACE600 system. For cryo-SEM, an FEI Helios NanoLab 650 FIB/SEM system equipped with Quorum PP3010T cryo system was used. Sample imaging temperature was -140°C, and sublimation was -70°C for 10 min. SEM images were analyzed using ImageJ (Abramoff et al. 2004).

### Stress testing

As a metric of stress-induced nanoemulsion instability, we calculated % change in nanoemulsion droplet sizes ( $d_z$ ) as follows:

$$\% \Delta d_z = \left( \frac{d_{z,f} - d_{z,i}}{d_{z,i}} \right) \times 100 \quad (2)$$

where  $d_{z,f}$  is the z-average nanoemulsion droplet size after the application of the stressor, and  $d_{z,i}$  is initial value for  $d_z$ .

The same quantity for variations in  $\zeta$  potential values is defined as follows:

$$\% \Delta \zeta = \left( \frac{\zeta_f - \zeta_i}{\zeta_i} \right) \times 100 \quad (3)$$

where  $\zeta_f$  is the  $\zeta$  potential of the nanoemulsion after the application of the stressor, and  $\zeta_i$  is initial value. For the purposes of this study, negative values of  $\% \Delta \zeta$  indicate destabilization by shifting of the  $\zeta$  potential value towards zero from either direction, while a positive  $\% \Delta \zeta$  indicates a shift to numerically greater  $\zeta$  potential values with attendant increase in interdroplet repulsions.

#### **Long-term storage**

To examine the effect of long term storage on the colloidal and chemical stability of the nanoemulsion, we stored them in tightly capped amber glass vials in a refrigerator at 4°C. Aliquots were periodically removed from the vials and diluted prior to DLS measurements. Measurements were performed immediately after high pressure homogenization, and after that, once every seven days for up to six weeks.

#### **Flash heating**

1 g of the optimized nanoemulsion was placed in a pre-heated water bath and the internal temperature of the nanoemulsion was maintained at 80°C for 1 minute. This protocol is a more extreme version of the high-temperature short-time (HTST) pasteurization process (typically, 71.5°C for 15 s) that fruit juices and milk beverages are subjected to in the industry (Negiz et al. 1998). The nanoemulsion was then allowed to cool to room temperature, and a part of it was diluted for DLS study. The rest was retained for cannabis potency evaluation.

#### **Freeze-thaw cycle**

1 g of the optimized nanoemulsion was placed in a freezer at a temperature of -20°C for 1 hour; then, the nanoemulsion was removed from the freezer and allowed to revert to room temperature. A part of the thawed nanoemulsion was diluted for DLS study, and the rest was retained for cannabis potency evaluation.

#### **Dilution and carbonation**

In a representative study, 1 mL of a representative cannabis nanoemulsion containing *ca.* 10 mg CBD was diluted to 355 mL (volume of a standard beverage can) using HPLC-grade water, thereby setting the CBD content at 10 mg per packaging unit. The diluted emulsion was then subjected to dynamic light scattering measurements. Given the absence of any stressors that might lead to cannabis decomposition or physical separation during this procedure, a potency evaluation wasn't performed. It is to be noted that the diluted nanoemulsion was almost transparent, which is a property sought by manufacturers

of cannabis-enriched carbonated-water type beverages (Donsi 2018; Saari and Chua 2020).

For the carbonation study, a similar protocol was adopted; however, for the dilution of 1 mL nanoemulsion concentrate to 355 mL, carbonated water obtained from a domestic Sodastream® unit operating in the “medium level of carbonation” mode was used. The resultant diluted beverage, containing approximately 10 mg of CBD, had a pH of 3.8.

#### **Change in pH**

1 g aliquots of the optimized CD<sub>CBD</sub> nanoemulsion were withdrawn at a starting pH of 3.6, and the pH of these aliquots were adjusted by the addition of 0.25 or 0.025(M) HCl or NaOH solution to values ranging from 1.9 to 9.4. The final weights of all aliquots were adjusted to the same value with HPLC water. After an incubation period of 12–18 h, these were diluted with solutions of matched pH values and subjected to DLS studies for measurement of average droplet diameters and zeta potentials. Meanwhile, cannabis potency evaluations were performed on the undiluted pH-adjusted aliquots.

#### **Ionic and covalent additives**

In a representative experiment, varying masses of CaCl<sub>2</sub> · 2 H<sub>2</sub>O were added to 5 mL portions of the optimized nanoemulsion so as to span a salt concentration range between 1 and 250 mM. To ensure complete dissolution of the salt, the aliquots were vortexed for 1 min each. After an incubation period of 12–18 h, it was noted that in vials with [CaCl<sub>2</sub> · 2 H<sub>2</sub>O] ≥ 100 mM, the nanoemulsions had partially or totally phase-separated (Figure S3). In order to perform the DLS studies, the aliquots were vortexed again for 1 min prior to dilution with calcium chloride solutions.

A similar protocol was followed to observe the impact of the addition of model non-ionic additive (here, sucrose, a sweetener) to a CD<sub>CBD</sub> nanoemulsion. To 5 mL portions of the optimized nanoemulsion were added varying masses of sucrose so as to span a sucrose concentration range between 1 and 500 mM (for reference, a 355 mL can of Coca-Cola typically has 39 g of sugar (The Coca-Cola Company 2021), which amounts to a sugar concentration of *ca.* 320 mM, assuming all the sugar is sucrose). To ensure complete dissolution of the sucrose, the aliquots were vortexed for 1 min each. After an incubation period of 12–18 h, 1 mL of aliquot was withdrawn from each sample in preparation for DLS evaluation. It is to be noted that cannabis potency studies were not carried out in the context of salt and sugar addition to the nanoemulsions, given that neither is expected to degrade cannabinoids and any cannabis potency loss from the bulk of the solution may be attributed to phase separation rather than to chemical transformations.

### Quantification of cannabis by HPLC

Quantitative cannabinoid analysis was performed at RPC, New Brunswick's provincial research organization (PRO), a research and technology organization (RTO) offering contract research and technical services in Fredericton. Cannabinoids were analyzed using an Agilent 1200 series HPLC system. Separation of cannabinoids was performed with a Phenomenex C18 column (Luna 5  $\mu\text{m}$ , 5.6 mm I.D. X 150 mm). The mobile phase was acetonitrile (A) and 25 mM ammonium formate solution (pH 3.75) (B) at a flow rate of 1 ml/min. The isocratic flow was 80% (A), 20% (B). The diode array detector was set at 228 nm. All solvents used were HPLC-grade and filtered via 0.20- $\mu\text{m}$  filters.

For calibration purposes, the cannabinoid standards (*vide supra*) were mixed at concentrations of 1 ppm, 2 ppm, 10 ppm, 20 ppm and 100 ppm. For analysis, 100 mg of sample was extracted with 30 mL of 90:10 methanol:chloroform, vortexed for 20 seconds and placed in an ultrasonic bath for 20 minutes. Afterwards, sample was centrifuged at 1200 rpm for 5 minutes. Supernatant was filtered using a 0.20  $\mu\text{m}$  filter, and fed into the HPLC.

### Statistical analysis


All DLS measurements were carried out in triplicate on at least two samples. Mean and standard deviations were calculated using Microsoft's Excel spreadsheet package (2016). For zeta potentials, the error bars represent the zeta deviation values obtained directly from the instrument.

## Results and discussion


### Optimization of emulsifier wt%

In nanoemulsions, the use of the least possible amount of emulsifier necessary to create a stable emulsion is recommended (*vide supra*). This is one of the primary justifications for the use of QNaturale<sup>®</sup> (15 wt% quillaja saponin) in our nanoemulsions: its low molecular mass and facile reduction of interfacial tension (*ca.* 5 mN/m for

MCT oil/water interface at 1 wt% quillaja saponin) leads to its rapid absorption at the lipid/water interface, and consequent stabilization of the resultant emulsions, even at low emulsifier concentrations (Chung et al. 2017). It is, however, of utmost importance to optimize the emulsifier wt%. We synthesized 10 wt% soybean oil nanoemulsions with 0.5, 1.5, 2.5, 5, 10, and 15 wt% of QNaturale<sup>®</sup>, and after 10 homogenization cycles at 30 Kpsi, observed the lipid phase droplet sizes and  $\zeta$  potentials (Fig. 2a). The difference between the average diameters of the lipid phase droplets of a nanoemulsion containing 0.5 wt% emulsifier and one containing 1.5 wt% emulsifier was about 125 nm. In contrast, when the emulsifier concentration was varied from 5 wt% to 10 wt%, this difference was only about 12 nm. Upon further increasing the QNaturale<sup>®</sup> concentration to 15 wt%, the emulsion clearly enters an 'emulsifier-rich region', and the droplet sizes become independent of the emulsifier concentration (McClements 2015). It is to be noted that the droplet sizes recorded by us (*ca.* 120 nm) matches well with the values recorded by Leibtag et al. for their olive oil/CBD/quillaja saponin nanoemulsion produced by a sonication method (*ca.* 110 nm), indicating a lower bound for quillaja saponin stabilized CBD/carrier oil droplets in the aqueous phase of a nanoemulsion, regardless of the emulsion preparation method (high pressure homogenization versus ultrasonic cavitation) or carrier oil type (soybean oil versus olive oil) (Leibtag and Peshkovsky 2020). It was suggested that this lower bound was a function of the character and packing capabilities of the surfactants. Similarly, more negative  $\zeta$  potential values were recorded at intermediate QNaturale<sup>®</sup> concentrations. The samples also showed enhanced monodispersity at greater emulsifier concentrations (Fig. 2b). These quantities, upon being remeasured after a week of storage under ambient conditions, did not exhibit any major deviations from their original values, other than demonstrating a smaller dependence of the  $\zeta$  potential on the QNE wt%. For further experiments, we



**Fig. 2** Variation of **a**  $d_z$ , **b** PDI, and **c**  $\zeta$  potential of a 10 wt% soybean oil nanoemulsion, immediately upon fabrication, and after a week, as a function of the wt% of QNaturale<sup>®</sup> (QNE) in the nanoemulsion




selected a QNaturale<sup>®</sup> concentration of 10 wt%, which roughly equated to a quillaja saponin concentration of approximately 1.5 wt% in the resultant nanoemulsion.

#### Optimization of number of homogenization cycles

Before the introduction of the  $CD_{CBD}$  within the nanoemulsion, we attempted to optimize factors such as the number of homogenization cycles, too few of which would lead to larger droplet sizes with attendant instability, while an excessive number would not only be operationally inefficient and expensive, but might also lead to emulsifier degradation and reduced colloidal stability of the emulsions (Peng et al. 2015). Therefore, we carried out 10 consecutive homogenization cycles at 30 Kpsi on a 10 wt% soybean oil nanoemulsion containing 10 wt% QNaturale<sup>®</sup> (Fig. 3), sampling aliquots after each processing cycle. There was a large reduction in soybean oil droplet sizes after the initial couple of homogenization cycles -  $d_z$  values changed from *ca.* 450 nm for the coarse emulsion, to 185 nm after one homogenization cycle, and to 165 nm after the second homogenization cycle. After this, the size largely stabilized; in fact, droplet size differences were within the limits of experimental error after 8–9 homogenization cycles. Therefore, we selected 10 as the optimal number of homogenization cycles for the generation of stable nanoemulsions. The entire high-pressure homogenization process for a coarse emulsion weighing *ca.* 40 g took 9.6 minutes for 10 homogenization cycles.  $\zeta$  potential values of the droplets showed a slight reduction (*ca.* 10 mV) over this range, possibly owing to detachment of charged moieties from the structure of quillaja saponin, caused either by the homogenization itself, or by the resultant localized heating within the emulsion during the process (Rampon et al. 2003; Flourey et al. 2003). This experiment was repeated with the optimized cannabis nanoemulsion (*vide infra*) in order to evaluate the impact of the number of HPH cycles on cannabis

potency (Fig. 4), with the measured weight of the CBD used in the emulsion preparation taken as 100%. Considering the CBD concentration in the nanoemulsion as a metric of cannabis potency, we found that the CBD concentration did not change beyond the limits of experimental error after the first two cycles, thereby indicating that CBD decomposition over 10 homogenization cycles is not a concern (Fig. 4). Incomplete mixing of the two phases and/or slight phase separation upon storage leading to heterogeneity in the samples explained the lower [CBD] values in the samples corresponding to the first two homogenization cycles analyzed by HPLC. Variation of



**Table 1**  $\zeta$  potential values of  $CD_{CBD}$  nanoemulsions with varying lipid phase compositions

Day	$CD_{CBD}$ % <sup>a</sup>	$\zeta$ (mV)	$\zeta$ deviation (mV)
1	10	-33.6	7.6
1	75	-28.1	5.7
8	10	-30.6	7.1
8	75	-21.1	7.4

<sup>a</sup>Percentage  $CD_{CBD}$  in the lipid phase; soybean oil is the other lipid phase component

the optimized nanoemulsion droplet sizes with the number of homogenization cycles were comparable to those recorded for the cannabis-free nanoemulsion (Figure S2).

### Optimization of lipid phase composition

In preliminary experiments, we noticed that cannabis-heavy lipid phases were more viscous and in general more difficult to handle. Moreover, nanoemulsions made with cannabis-majority lipid phases had broad, bimodal droplet size distributions and tended to exhibit droplet agglomeration upon prolonged storage. Therefore, we fabricated a series of lipid phases with varying  $CD_{CBD}$  contents (100, 75, 50, 25, 16.6, and 10 wt% of  $CD_{CBD}$ ) mixed with appropriate amounts of soy oil. These were then used to generate 10 wt% lipid phase nanoemulsions containing 10 wt% QNaturale<sup>®</sup> as described before. Dispersed phase droplet sizes, polydispersity index values, and  $\zeta$  potentials were measured for these systems both immediately after creation, as well as after 7 days under ambient storage conditions. It is evident that the average lipid phase droplet sizes decrease (Fig. 5a) and the dispersion becomes monodisperse (Fig. 5b, Figure S2) as the cannabis oil wt% is reduced, potentially owing to the higher viscosity of the cannabis extract compared to soybean oil. The  $\zeta$  potentials, on the other hand, remain relatively unchanged within the limits of experimental error, both as a function

of the lipid phase composition, as well as upon prolonged storage (Table 1). Further experiments were conducted with the lipid phase containing 16.6 wt% of  $CD_{CBD}$ .


### Imaging the optimized cannabis/soybean oil-in-water nanoemulsion

Representative SEMs of the optimized  $CD_{CBD}$  nanoemulsion may be found in Fig. 6. The size distribution, calculated from data obtained by measuring the diameters of >250 droplets, revealed the average droplet size to be 115 nm (SD= 39 nm), which is somewhat smaller than the DLS size data. This, however, is not unexpected, since DLS tends to 'over-weigh' larger particles owing to their greater light-scattering ability (Souza et al. 2016). Moreover, DLS considers a layer of water molecules surrounding the lipid droplet, generating what is called the hydrodynamic radius, which is inevitably greater than the actual radius of the lipid droplet.


In the presence of 50 mM  $CaCl_2 \cdot 2 H_2O$ , the nanoemulsion droplets show an increase in size, with the average droplet diameter increasing to 165 nm (SD= 48 nm); this trend is in agreement with the DLS data. Optical micrographs of the optimized nanoemulsion with and without additives may be found in Figure S5 as well as insets in the appropriate graphs.

### Impact of physical stressors

The impact of physical stressors on the stability and potency of cannabis in the optimized cannabis nanoemulsion have been summarized in Table 2, with the  $d_{z,f}$  and  $\zeta_f$  values obtained from measurement of control nanoemulsion samples *sans* exposure to stressors. On the basis of the  $\% \Delta d_z$  values, we conclude that pasteurization-style flash heating and a single freeze-thaw cycle at  $-20^\circ C$  has minimal impact on the nanoemulsion parameters except a slight increase in droplet sizes. Dilution to 355 mL, using either still or carbonated water, reduces average



**Fig. 5** Variation of **a**  $d_z$  and **b** PDI of a phytocannabinoid nanoemulsion, immediately upon fabrication, and after a week, as a function of the wt% of  $CD_{CBD}$  in the lipid phase of the nanoemulsion



**Fig. 6** **a** SEM of the optimized nanoemulsion, showing lipid phase droplets; (inset) close-up of the nanoemulsion droplets; **b** SEM of the nanoemulsion in the presence of 50 mM  $\text{CaCl}_2$ . Size distribution profiles are underneath the respective SEMs

droplet size slightly, while droplet size distribution profiles remain unaltered, which confirms that our nanoemulsion is kinetically metastable. However, upon storage of the nanoemulsion for 6 weeks under ambient conditions, slight 'ringing' (creation of a very thin whitish creamy ring on the glass surface at the top of the emulsion) (Kumar and Sarkar 2018) was observed, likely due to difference in density between the oil and water phases, and the  $\zeta$  potential values fell. Concentration of CBD in the optimized

nanoemulsion decreased by *ca.* 17% after 6 weeks of storage upon comparison to the initial concentration of CBD in the freshly synthesized nanoemulsion (Fig. 7).

#### Impact of chemical stressors

In practical applications, cannabinoid nanoemulsions are expected to retain their desirable emulsion properties over a wide range of pH values and ionic strengths, depending upon product composition. Factors such as

**Table 2** Impact of stressors on optimized  $\text{CD}_{\text{CBD}}$  nanoemulsion parameters


System	Stressor	$d_{z,f}$ (nm)	$\% \Delta d_z$	$\zeta_f$ (mV)	$\% \Delta \zeta_f$	[CBD] ( $\text{mg}\cdot\text{g}^{-1}$ )
1 <sup>a</sup>	none	125.5	N/A	-32.2	N/A	10.8
2	heat	136.8	9	-30.9	-4	10.4
3	freeze-thaw	142.5	13.5	-30.2	-6.2	10.6
4	dilution <sup>b</sup>	123.6	-1.5	-25.6	-19.7	N/A <sup>c</sup>
5	dilution/carbonation <sup>d</sup>	135.8	-3.5	-25.0	-8.9	N/A <sup>c</sup>
6	6 weeks storage	130.0	3.6	-29.4	-8.7	9.1

<sup>a</sup>Optimized nanoemulsion, as synthesized

<sup>b</sup>1 mL nanoemulsion added to HPLC water to obtain a final volume of 355 mL

<sup>c</sup>[CBD] not measured as dilution is not expected to change it

<sup>d</sup>1 mL nanoemulsion added to carbonated water to obtain a final volume of 355 mL




pH, the addition of ionic salts, and organic additives such as sugars can have profound effects on the stability of nanoemulsions (McClements 2004).

#### Effect of pH

Changes in pH can protonate or deprotonate terminal functional groups of emulsifiers, thereby modulating interdroplet electrostatic repulsive forces, and ultimately affecting droplet coalescence rates (Chung et al. 2017). For our optimized nanoemulsion, however, a moderate increase in droplet size (*ca.* 30 nm) was observed upon lowering the pH to highly acidic regime (pH < 2), while no changes in droplet sizes were recorded upon raising the pH to 9 (Fig. 8a). Visual observation of the emulsions indicated that creaming occurred at pH 2 and below, which is consistent with previous reports (Yang et al. 2013); also, optical microscopy of the optimized nanoemulsion at pH 1.6 revealed the presence of extensive flocs. These results seem to indicate that despite minimal particle coalescence, the emulsions show a high degree of flocculation at pH 2 and below. Protonation of the terminal  $-\text{COO}^-$

groups of quillaja saponin leading to reduced interdroplet repulsion at very low pH values may be invoked to explain this observation. As expected, zeta potentials became steadily more negative with increase in pH. These results indicate that  $\text{CD}_{\text{CBD}}$  nanoemulsions stabilized by quillaja saponin are stable across most of the pH range present within food products (pH 3–8), but very acidic matrices



might lead to flocculation in such nanoemulsions. Quantitative analysis of emulsions at various pH values indicated a steady concentration of CBD that didn't change at different pH values, even after a month of incubation in the fridge at 4°C, suggesting that flocculation in this case is not accompanied by irreversible phase separation (Figure S6).

#### **Effect of covalent additives**

The addition of varying amounts of sucrose, a commonly used sweetener, initially caused no changes to the  $CD_{CBD}$  nanoemulsion droplet sizes and  $\zeta$  potential values; however, lipid droplet sizes begin to increase when the sucrose concentration exceeds 250 mM (Fig. 8b). At 500 mM sucrose concentrations, the average droplet sizes of the optimized  $CD_{CBD}$  nanoemulsion increased from *ca.* 130 nm to *ca.* 170 nm. From cryo-SEM, average lipid phase droplet size of the optimized nanoemulsion in the presence of *ca.* 100 mM sucrose was found to be  $182.3 \pm 48.4$  nm, which is in rough agreement with DLS size data (Figure S4). This observation is explained by the growing disparity between the density and viscosity of the aqueous and lipid phases as more and more sucrose is dissolved in the aqueous phase (McClements 2015). The  $\zeta$  potential values of the nanoemulsion showed a slight increase (*ca.* 10 mV) with increase in sucrose concentration; this was seen to be reproducible over multiple experiments. This was likely brought about by the expanding location of the slipping plane during electrophoretic movement with increasing concentration of sugars, due to the formation of a viscous hydration layer on the surface of the nanodroplets, as recorded previously by Matsumoto (1994). No noticeable changes in the polydispersity index of the samples were noticed even at the highest sucrose concentrations.

#### **Effect of ionic additives**


The addition of  $\text{CaCl}_2 \cdot 2 \text{H}_2\text{O}$  was intended to mimic the effect of hard water; also, essential minerals are often intentionally added to nutraceutical products (Keownaneechai and McClements 2002). It is already known that the addition of salts such as  $\text{CaCl}_2 \cdot 2 \text{H}_2\text{O}$  can lead to instability and phase separation in nanoemulsions containing quillaja saponin; nevertheless, it is superior in this context to other emulsifiers such as beet pectins, given that it offers both electrostatic and steric stability to the lipid phase droplets (Ralla et al. 2017). Previous studies have demonstrated the quillaja saponin stabilized nanoemulsions showed droplet aggregation and flocculation at  $[\text{NaCl}] \geq 300$  mM (Ralla et al. 2017). In our study, we found that  $CD_{CBD}$  nanoemulsion droplet sizes remained in the sub- $\mu\text{m}$  region despite some increase in average diameters at  $[\text{CaCl}_2 \cdot 2 \text{H}_2\text{O}] \leq 50$  mM (Fig. 6b); at 100 mM, slow but discernible phase separation was initiated,

while above 100 mM rapid separation of nanoemulsion into serum layer and creamy layer could be observed (Figure S3). The primary reason for this observation is the high screening capacity of the  $\text{Ca}^{2+}$  ion, which reduces the interdroplet Debye screening length, and consequently interdroplet repulsion, until attractive interactions between droplets (such as van der Waal forces) predominate, leading to droplet coalescence and/or flocculation (McClements 2015). This effect is supposed to be especially strong in emulsions with  $\zeta$ -potentials with magnitudes above 25 mV (as is the case here) in the presence of polyvalent cations (McClements 2015; Ralla et al. 2017) (Fig. 9a). We attempted to measure the  $\zeta$ -potentials of our  $\text{CaCl}_2$ -containing nanoemulsion aliquots (Fig. 9b) but the high ionic strengths led to large errors in measured  $\zeta$ -potential values, possibly owing to effects such as ion-droplet and droplet-ion-droplet bridging (Dickinson 2010). It is clear, however, that drastic reduction of  $\zeta$ -potential could be observed with increasing  $[\text{CaCl}_2 \cdot 2 \text{H}_2\text{O}]$ , with values dropping to zero between 150 and 200 mM  $[\text{CaCl}_2 \cdot 2 \text{H}_2\text{O}]$  (Fig. 9b). In order to visualize our optimized nanoemulsion in the presence of various additives, optical microscopy was carried out, and the respective micrographs may be found in the SI (Figure S5).

The relatively high stability of the optimized  $CD_{CBD}$  nanoemulsion may be attributed to the fact that the creaming rate is directly proportional to the square of the droplet diameter - an emulsion with nanosized droplets, therefore, has a reduced rate of gravitational separation (McClements 2015). Additionally, Brownian motion effects that favor a homogeneous distribution of droplets throughout the system counteract the gravitational forces when the droplets are sufficiently small, as is the case for our optimized nanoemulsion. The charged  $-\text{COO}^-$  termini present in the quillaja saponin structure also offers electrostatic stabilization by enhancing interdroplet repulsion. With respect to their chemical stability, the documented high free radical quenching capacity of quillaja saponin may help minimize oxidative degradation of  $CD_{CBD}$  within the emulsion lipid phase droplets (Uluata et al. 2016; Toppel et al. 2017).

#### **Conclusions**

This report presents the design, creation, and stress-testing results for a novel cannabidiol-enriched nanoemulsion as a representative cannabis-containing nutraceutical suitable for ingestion. After optimization of the number of high pressure homogenization cycles, emulsifier wt%, and lipid phase composition, we produced a CBD-enriched soybean oil nanoemulsion in water in the presence of 1.5 wt% quillaja saponin with average lipid phase droplet sizes of *ca.* 120 nm and  $\zeta$  potential values of approx. -30 mV. The nanoemulsion proved to be stable over a period of 6 weeks, with minor



creaming, and was resistant to potential instability owing to flash-heating, cooling, dilution and addition of carbonated water. Droplet agglomeration became a concern, however, at  $pH < 2$ , as well in the presence of excessive amounts of additives ( $[sucrose] \geq 250$  mM,  $[CaCl_2] > 100$

mM). Cannabis potency, measured in terms of CBD concentration, changed by  $\sim 17\%$  over a period of 6 weeks, and was largely unaffected by other examined stressors. It is expected that this report will be the first of a series stemming from a detailed study of various cannabinoid nanoemulsions suitable for the nutraceutical industry in Canada and other countries where the therapeutic potentials of cannabis are finally being widely acknowledged and commercialized.

#### Abbreviations

CBD: Cannabidiol, IUPAC name: 2-[(1R,6R)-3-methyl-6-prop-1-en-2-ylcyclohex-2-en-1-yl]-5-pentylbenzene-1,3-diol;  $CD_{CBD}$ : CBD-enriched cannabis distillate;  $CaCl_2 \cdot 2 H_2O$ : Calcium chloride dihydrate; DLS: Dynamic light scattering; HPLC: High-performance liquid chromatography; PDI: Polydispersity index; SEM: Scanning electron microscopy; THC:  $\Delta^9$ -tetrahydrocannabinol, IUPAC Name: (6aR,10aR)-6,6,9-trimethyl-3-pentyl-6a,7,8,10a-tetrahydrobenzo [c]chromen-1-ol

#### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s42238-021-00094-w>.

**Additional file 1:** See additional file named **SI.pdf** for quantitative analysis of the CBD-enriched cannabis distillate used in this study, droplet size distribution profiles for  $CD_{CBD}$  nanoemulsions at various lipid phase compositions, variation of the droplet sizes of the optimized nanoemulsion as a function of the number of homogenization cycles, photographic representation of nanoemulsion destabilization and phase separation in the presence of added  $CaCl_2$ , cryo-SEM of the optimized nanoemulsion in the presence of 100 mM sucrose, and data for the variation of  $CD_{CBD}$  concentration in the optimized nanoemulsion as a function of pH.

#### Acknowledgements

We thank Troy Smith and his team at RPC for the HPLC analysis of cannabinoids in our systems. SEM and cryo-SEM imaging were performed at the 4D labs in Simon Fraser University, BC. We also thank Prof. Holger Eichhorn for the use of the optical microscope. The TOC figure was created with BioRender.

#### Disclaimer

The opinions expressed in this manuscript are those of the authors and do not represent statements of position, intention or strategy of Peak Processing Solutions.

#### Authors' contributions

Conceptualization: AB, JB, JFT; Formal analysis: AB; Funding acquisition: JFT, JB; Investigation: AB, RS; Methodology: AB, JFT; Resources: JFT, JB; Supervision: JFT; Visualization: AB; Writing - original draft: AB; Writing - review and editing: All authors. All authors read and approved the final manuscript.

#### Funding

We thank the MITACS Accelerate Fellowship program (MITACS IT16195 to JFT for salary to AB), the University of Windsor (Ignite funding to RS), the Natural Sciences and Engineering Research Council of Canada (2018-06338 to JFT) and Peak Processing for supporting this research.

#### Availability of data and materials

The authors maintain all raw data for a minimum of 15 years and this is available from the senior corresponding author upon reasonable request. Additional data is provided as supplementary information in the accompanying files.

#### Declarations

#### Ethical approval and consent to participate

Not applicable to this study.

### Consent for publication

All the authors declare that they have reviewed and approved the final manuscript prior to communication and have no objections to the publication of the same.

### Competing interests

AB is funded by a MITACS Accelerate Fellowship with Peak Processing as the industrial partner. JB is the manager of scientific development at Peak Processing.

### Author details

<sup>1</sup>Department of Chemistry and Biochemistry, University of Windsor, 401 Sunset Ave., Windsor, ON, Canada. <sup>2</sup>Peak Processing Solutions, Tecumseh, ON, Canada.

Received: 2 November 2020 Accepted: 5 August 2021

Published online: 23 September 2021

### References

- Abioye A, Ayodele O, Marinkovic A, Patidar R, Akinwekomi A, Sanyaolu A.  $\delta$ 9-tetrahydrocannabinol (thcv): a commentary on potential therapeutic benefit for the management of obesity and diabetes. *J Cannabis Res.* 2020;2(1):1–6.
- Abramoff MD, Magalhaes PJ, Ram SJ. Image processing with imagej. *Biophoton Int.* 2004;11(7):36–42.
- Acosta E. Bioavailability of nanoparticles in nutrient and nutraceutical delivery. *Curr Opin Colloid Interf Sci.* 2009;14(1):3–15.
- Atsmon J, Cherniakov I, Izgelov D, Hoffman A, Domb AJ, Deutsch L, Deutsch F, Heffetz D, Sacks H. Pt1401, a new formulation based on pro-nano dispersion technology, improves oral cannabinoids bioavailability in healthy volunteers. *J Pharm Sci.* 2018;107(5):1423–9.
- Bai L, Huan S, Gu J, McClements DJ. Fabrication of oil-in-water nanoemulsions by dual-channel microfluidization using natural emulsifiers: Saponins, phospholipids, proteins, and polysaccharides. *Food Hydrocoll.* 2016;61:703–11.
- Cameron EC, Hemingway SL. Cannabinoids for fibromyalgia pain: a critical review of recent studies (2015–2019). *J Cannabis Res.* 2020;2:1–11.
- Chen L, Subirade M. Elaboration and characterization of soy/zein protein microspheres for controlled nutraceutical delivery. *Biomacromolecules.* 2009;10(12):3327–34.
- Cherniakov I, Izgelov D, Domb AJ, Hoffman A. The effect of pro nanolipospheres (pnl) formulation containing natural absorption enhancers on the oral bioavailability of delta-9-tetrahydrocannabinol (thc) and cannabidiol (cbd) in a rat model. *Eur J Pharm Sci.* 2017;109:21–30.
- Chung H, Kim TW, Kwon M, Kwon IC, Jeong SY. Oil components modulate physical characteristics and function of the natural oil emulsions as drug or gene delivery system. *J Control Release.* 2001;71(3):339–50.
- Chung C, Sher A, Rousset P, McClements DJ. Influence of homogenization on physical properties of model coffee creamers stabilized by quillaja saponin. *Food Res Int.* 2017;99:770–7.
- Government of Canada. Department of Justice, Government of Canada. Cannabis Legalization and Regulation. 2018. <https://www.justice.gc.ca/eng/cj-jp/cannabis/>. Accessed 2 Sept 2021.
- Dickinson E. Food emulsions and foams: Stabilization by particles. *Curr Opin Colloid Interf Sci.* 2010;15(1-2):40–9.
- Donsi F. Applications of nanoemulsions in foods. In: *Nanoemulsions*. Amsterdam: Academic Press, Elsevier; 2018. p. 349–77. <https://doi.org/10.1016/b978-0-12-811838-2.00011-4>.
- Elgart A, Cherniakov I, Aldouby Y, Domb AJ, Hoffman A. Lipospheres and pro-nano lipospheres for delivery of poorly water soluble compounds. *Chem Phys Lipids.* 2012;165(4):438–53.
- Floury J, Desrumaux A, Axelos MAV, Legrand J. Effect of high pressure homogenisation on methylcellulose as food emulsifier. *J Food Eng.* 2003;58(3):227–38.
- Grotenhermen F. Pharmacokinetics and pharmacodynamics of cannabinoids. *Clin Pharmacokinet.* 2003;42(4):327–60.
- Hanus LO, Meyer SM, Munoz E, Tagliatalata-Scafati O, Appendino G. Phytocannabinoids: a unified critical inventory. *Nat Prod Rep.* 2016;33(12):1357–92.
- Hsu J-P, Nacu A. Behavior of soybean oil-in-water emulsion stabilized by nonionic surfactant. *J Colloid Interfac Sci.* 2003;259(2):374–81.
- Jaiswal M, Dudhe R, Sharma P. Nanoemulsion: an advanced mode of drug delivery system. *3 Biotech.* 2015;5(2):123–7.
- Keowmaneechai E, McClements DJ. Effect of cacl2 and kcl on physicochemical properties of model nutritional beverages based on whey protein stabilized oil-in-water emulsions. *J Food Sci.* 2002;67(2):665–71.
- Khan R, Naveed S, Mian N, Fida A, Raafey MA, Aedma KK. The therapeutic role of cannabidiol in mental health: a systematic review. *J Cannabis Res.* 2020;2(1):1–21.
- Kharat M, McClements DJ. Fabrication and characterization of nanostructured lipid carriers (nlc) using a plant-based emulsifier: Quillaja saponin. *Food Res Int.* 2019;126:108601.
- Knaub K, Sartorius T, Dharsono T, Wacker R, Wilhelm M, Schoen C. A novel self-emulsifying drug delivery system (sedds) based on vesisorb® formulation technology improving the oral bioavailability of cannabidiol in healthy subjects. *Molecules.* 2019;24(16):2967.
- Kumar DL, Sarkar P. Encapsulation of bioactive compounds using nanoemulsions. *Environ Chem Lett.* 2018;16(1):59–70.
- Leibtag S, Peshkovsky A. Cannabis extract nanoemulsions produced by high-intensity ultrasound: Formulation development and scale-up. *J Drug Deliv Sci Technol.* 2020;60:101953. <https://doi.org/10.1016/j.jddst.2020.101953>.
- Marangoni IP, Marangoni AG. Cannabis edibles: dosing, encapsulation, and stability considerations. *Curr Opin Food Sci.* 2019;28:1–6.
- Maroon J, Bost J. Review of the neurological benefits of phytocannabinoids. *Surg Neurol Int.* 2018;9:91(1)–91(13).
- Matsumoto S. Proteins and sugars affecting the zeta potential and stability of dispersed vesicular globules in w/o/w emulsions. In: *Food hydrocolloids*. Springer; 1994. p. 399–408. [https://doi.org/10.1007/978-1-4615-2486-1\\_62](https://doi.org/10.1007/978-1-4615-2486-1_62).
- McClements DJ. Protein-stabilized emulsions. *Curr Opin Colloid Interf Sci.* 2004;9(5):305–13.
- McClements DJ. *Food Emulsions: Principles, Practices, and Techniques*. Boca Raton: CRC press; 2015.
- McClements DJ. Enhancing efficacy, performance, and reliability of cannabis edibles: Insights from lipid bioavailability studies. *Annu Rev Food Sci Technol.* 2020;11:45–70.
- Mclver B. Cannabis Legalization World Map. 2017. <https://www.cannabisbusinesstimes.com/article/cannabis-legalization-world-map/>. Accessed 14 Sep 2021.
- Mouritsen O, Styrbaek K. *Mouthfeel: How Texture Makes Taste*. New York: Columbia University Press; 2017.
- Nakano Y, Tajima M, Sugiyama E, Sato VH, Sato H. Development of a novel nanoemulsion formulation to improve intestinal absorption of cannabidiol. *Med Cannabis Cannabinoids.* 2019;2(1):35–42.
- Negiz A, Ramanauskas P, Cinar A, Schlessler JE, Armstrong DJ. Modeling, monitoring and control strategies for high temperature short time pasteurization systems—1. empirical model development. *Food Control.* 1998;9(1):1–15.
- Panghal A, Chhikara N, Anshid V, Charan MVS, Surendran V, Malik A, Dhull SB. Nanoemulsions: A promising tool for dairy sector. In: *Nanobiotechnology in Bioformulations*. Basel: Springer International Publishing; 2019. p. 99–117. [https://doi.org/10.1007/978-3-030-17061-5\\_4](https://doi.org/10.1007/978-3-030-17061-5_4).
- Parker LA, Mechoulam R, Schlievert C. Cannabidiol, a non-psychoactive component of cannabis and its synthetic dimethylheptyl homolog suppress nausea in an experimental model with rats. *Neuroreport.* 2002;13(5):567–70.
- Peng J, Dong W-J, Li L, Xu J-M, Jin D-J, Xia X-J, Liu Y-L. Effect of high-pressure homogenization preparation on mean globule size and large-diameter tail of oil-in-water injectable emulsions. *J Food Drug Anal.* 2015;23(4):828–35.
- Raikos V, Ranawana V. Designing emulsion droplets of foods and beverages to enhance delivery of lipophilic bioactive components—a review of recent advances. *Int J Food Sci Technol.* 2017;52(1):68–80.
- Ralla T, Salminen H, Edelmann M, Dawid C, Hofmann T, Weiss J. Stability of emulsions using a new natural emulsifier: sugar beet extract (beta vulgaris l). *Food Biophys.* 2017;12(3):269–78.
- Rampon V, Riaublanc A, Anton M, Genot C, McClements DJ. Evidence that homogenization of bsa-stabilized hexadecane-in-water emulsions induces structure modification of the nonadsorbed protein. *J Agric Food Chem.* 2003;51(20):5900–5.
- Reekie TA, Scott MP, Kassiou M. The evolving science of phytocannabinoids. *Nat Rev Chem.* 2017;2(1):1–12.
- Reichert CL, Salminen H, Weiss J. Quillaja saponin characteristics and functional properties. *Annu Rev Food Sci Technol.* 2019;10(1):43–73.

- Saari NHM, Chua LS. Nano-based products in beverage industry. In: Nanoengineering in the Beverage Industry. Amsterdam: Academic Press, Elsevier; 2020. p. 405–36. <https://doi.org/10.1016/b978-0-12-816677-2.00014-4>.
- Scherma M, Masia P, Deidda M, Fratta W, Tanda G, Fadda P. New perspectives on the use of cannabis in the treatment of psychiatric disorders. *Medicines*. 2018;5(4):107.
- Shahbazi F, Grandi V, Banerjee A, Trant JF. Cannabinoids and cannabinoid receptors: The story so far. *iScience*. 2020;23(7):101301. <https://doi.org/10.1016/j.isci.2020.101301>.
- Solans C, Izquierdo P, Nolla J, Azemar N, Garcia-Celma MJ. Nano-emulsions. *Curr Opin Colloid Interf Sci*. 2005;10(3-4):102–10.
- Souza TGF, Ciminelli VST, Mohallem NDS. A comparison of TEM and DLS methods to characterize size distribution of ceramic nanoparticles. *J Phys Conf Ser*. 2016;733:012039.
- Sun Y, Xia Z, Zheng J, Qiu P, Zhang L, McClements DJ, Xiao H. Nanoemulsion-based delivery systems for nutraceuticals: Influence of carrier oil type on bioavailability of pterostilbene. *J Funct Foods*. 2015;13:61–70.
- Sze A, Erickson D, Ren L, Li D. Zeta-potential measurement using the smoluchowski equation and the slope of the current–time relationship in electroosmotic flow. *J Colloid Interf Sci*. 2003;261(2):402–10.
- Takino T, Konishi K, Takakura Y, Hashida M. Long circulating emulsion carrier systems for highly lipophilic drugs. *Biol Pharma Bull*. 1994;17(1):121–5.
- Teng Z, Luo Y, Wang Q. Nanoparticles synthesized from soy protein: preparation, characterization, and application for nutraceutical encapsulation. *J Agric Food Chem*. 2012;60(10):2712–20.
- The Coca-Cola Company. How Much Sugar Is in Coca-Cola? 2021. <https://www.coca-colacompany.com/faqs/how-much-sugar-is-in-coca-cola>. Accessed 2 Sept 2021.
- Tippel J, Gies K, Harbaum-Piayda B, Steffen-Heins A, Drusch S. Composition of quillaja saponin extract affects lipid oxidation in oil-in-water emulsions. *Food Chem*. 2017;221:386–94.
- Turner SE, Williams CM, Iversen L, Whalley BJ. Molecular pharmacology of phytocannabinoids. In: Kinghorn AD, Falk H, Gibbons S, Kobayashi J, editors. *Phytocannabinoids: Unraveling the Complex Chemistry and Pharmacology of Cannabis Sativa*. Basel: Springer International Publishing; 2017. p. 61–101.
- Uluata S, Decker EA, McClements DJ. Optimization of nanoemulsion fabrication using microfluidization: role of surfactant concentration on formation and stability. *Food Biophys*. 2016;11(1):52–9.
- Vasudevan K, Stahl V. Cannabinoids infused mouthwash products are as effective as chlorhexidine on inhibition of total-culturable bacterial content in dental plaque samples. *J Cannabis Res*. 2020;2(1):1–9.
- Yang Y, Leser ME, Sher AA, McClements DJ. Formation and stability of emulsions using a natural small molecule surfactant: Quillaja saponin (q-naturale®). *Food Hydrocoll*. 2013;30(2):589–96.
- Zhang J, Bing L, Reineccius GA. Comparison of modified starch and quillaja saponins in the formation and stabilization of flavor nanoemulsions. *Food Chem*. 2016;192:53–9.
- Zhu Z, Wen Y, Yi J, Cao Y, Liu F, McClements DJ. Comparison of natural and synthetic surfactants at forming and stabilizing nanoemulsions: Tea saponin, quillaja saponin, and tween 80. *J Colloid Interf Sci*. 2019;536:80–7.
- Zou S, Kumar U. Cannabinoid receptors and the endocannabinoid system: signaling and function in the central nervous system. *Int J Mol Sci*. 2018;19(3):833.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

